


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 405/00, A61K 31/557	A1	(11) International Publication Number: WO 95/19964 (43) International Publication Date: 27 July 1995 (27.07.95)
(21) International Application Number: PCT/US95/00288 (22) International Filing Date: 9 January 1995 (09.01.95) (30) Priority Data: 08/183,682 19 January 1994 (19.01.94) US (71) Applicant: ALLERGAN, INC. [US/US]; 2525 Dupont Drive, P.O. Box 19534, Irvine, CA 92713-9534 (US). (72) Inventor: WOODWARD, David, F.; 23152 Tulip Street, El Toro, CA 92630 (US).	(81) Designated States: AU, CA, CN, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: EP₂-RECEPTOR AGONISTS AS AGENTS FOR LOWERING INTRAOCULAR PRESSURE

(57) Abstract

The invention relates to (±) trans- 2-[-4(1-hydroxyhexyl) phenyl]-5-oxocyclopentaneheptanoic acid, and ester and unsaturated derivatives thereof, prostaglandin derivatives. More particularly, the present invention concerns compounds of formula (I), wherein the wavy bonds indicate the α or β configuration, the dashed bond represents either a single or double bond, which double bond may be a cis or trans double bond and R is a saturated or unsaturated acyclic hydrocarbon group having from 1 to about 20 carbon atoms, or -(CH₂)_mR₁ wherein m is 0-10, and R₁ is an aliphatic ring having from about 3 to about 7 carbon atoms, or an aryl or heteroaryl ring having from about 4 to about 10 carbon atoms and wherein the heteroatom is selected from the group consisting of N, O and S. The compounds of the present invention are potent ocular hypotensives, and are particularly suitable for the management of glaucoma.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

EP₂-RECEPTOR AGONISTS AS AGENTS FOR LOWERING INTRAOCULAR PRESSURE

5

Field of the Invention

The present invention relates to the use of EP₂ receptor agonists to lower the intraocular pressure of mammals and thus are useful in treating glaucoma. In particular, (±) trans-
10 2-[-4(1-hydroxyhexyl) phenyl]-5-oxocyclopentaneheptanoic acid, and ester and unsaturated derivatives thereof, are effective for the management of glaucoma.

15

Background of the Invention

Ocular hypotensive agents are useful in the treatment of a number of various ocular hypertensive conditions, such as post-surgical and post-laser trabeculectomy ocular hypertensive episodes, glaucoma, and as presurgical adjuncts.

20

Glaucoma is a disease of the eye characterized by increased intraocular pressure. On the basis of its etiology, glaucoma has been classified as primary or secondary. For example, primary glaucoma in adults (congenital glaucoma)
25 may be either open-angle or acute or chronic angle-closure. Secondary glaucoma results from pre-existing ocular diseases such as uveitis, intraocular tumor or an enlarged cataract.

30

The underlying causes of primary glaucoma are not yet known. The increased intraocular tension is due to the obstruction of aqueous humor outflow. In chronic open-angle glaucoma, the anterior chamber and its anatomic structures appear normal, but drainage of the aqueous humor is impeded. In acute or chronic angle-closure glaucoma, the anterior
35 chamber is shallow, the filtration angle is narrowed, and the iris may obstruct the trabecular meshwork at the entrance of the canal of Schlemm. Dilation of the pupil may push the root

of the iris forward against the angle, and may produce pupillary block and thus precipitate an acute attack. Eyes with narrow anterior chamber angles are predisposed to acute angle-closure glaucoma attacks of various degrees of severity.

5

Secondary glaucoma is caused by any interference with the flow of aqueous humor from the posterior chamber into the anterior chamber and subsequently, into the canal of Schlemm. Inflammatory disease of the anterior segment may prevent aqueous escape by causing complete posterior synechia in iris bombe and may plug the drainage channel with exudates. Other common causes are intraocular tumors, enlarged cataracts, central retinal vein occlusion, trauma to the eye, operative procedures and intraocular hemorrhage.

15

Considering all types together, glaucoma occurs in about 2% of all persons over the age of 40 and may be asymptotic for years before progressing to rapid loss of vision. In cases where surgery is not indicated, topical β -adrenoreceptor antagonists have traditionally been the drugs of choice for treating glaucoma.

20

Prostaglandins were earlier regarded as potent ocular hypertensives; however, evidence accumulated in the last two decades shows that some prostaglandins are highly effective ocular hypotensive agents and are ideally suited for the long-term medical management of glaucoma. (See, for example, Starr, M.S. Exp. Eye Res. 1971, 11, pp. 170-177; Bito, L. Z. Biological Protection with Prostaglandins Cohen, M. M., ed., Boca Raton, Fla, CRC Press Inc., 1985, pp. 231-252; and Bito, L. Z., Applied Pharmacology in the Medical Treatment of Glaucomas Drance, S. M. and Neufeld, A. H. eds., New York, Grune & Stratton, 1984, pp. 477-505). Such prostaglandins include $\text{PGF}_2\alpha$, $\text{PGF}_1\alpha$, PGE_2 , and certain lipid-soluble esters, such as C_1 to C_5 alkyl esters, e.g. 1-isopropyl ester, of such compounds.

25

30

35

In the United States Patent No. 4,599,353 certain prostaglandins, in particular PGE₂ and PGF₂ α and the C₁ to C₅ alkyl esters of the latter compound, were reported to possess ocular hypotensive activity and were recommended for use in glaucoma management.

Although the precise mechanism is not yet known, recent experimental results indicate that the prostaglandin-induced reduction in intraocular pressure results from increased uveoscleral outflow [Nilsson et al., Invest. Ophthalmol. Vis. Sci. 28(suppl), 284 (1987)].

The isopropyl ester of PGF₂ α has been shown to have significantly greater hypotensive potency than the parent compound, which was attributed to its more effective penetration through the cornea. In 1987 this compound was described as "the most potent ocular hypotensive agent ever reported." [See, for example, Bito, L. Z., Arch. Ophthalmol. 105, 1036 (1987), and Siebold et al., Prodrug 5, 3 (1989)].

Whereas prostaglandins appear to be devoid of significant intraocular side effects, ocular surface (conjunctival) hyperemia and foreign-body sensation have been consistently associated with the topical ocular use of such compounds, in particular PGF₂ α and its prodrugs, e.g. its 1-isopropyl ester, in humans. The clinical potential of prostaglandins in the management of conditions associated with increased ocular pressure, e.g. glaucoma, is greatly limited by these side effects.

Certain phenyl and phenoxy mono, tri and tetra nor prostaglandins and their 1-esters are disclosed in European Patent Application 0,364,417 as useful in the treatment of glaucoma or ocular hypertension.

In a series of co-pending United States patent applications assigned to Allergan, Inc. prostaglandin esters with increased ocular hypotensive activity accompanied with no or substantially reduced side-effects are disclosed. The co-pending USSN 386,835 (filed 27 July 1989), relates to certain 11-acyl-prostaglandins, such as 11-pivaloyl, 11-acetyl, 11-isobutyryl, 11-valeryl, and 11-isovaleryl PGF₂ α . Intraocular pressure reducing 15-acyl prostaglandins are disclosed in the co-pending application USSN 357,394 (filed 25 May 1989). Similarly, 11,15- 9,15- and 9,11-diesters of prostaglandins, for example 11,15-dipivaloyl PGF₂ α are known to have ocular hypotensive activity. See the co-pending patent applications USSN No. 385,645 filed 27 July 1990, now U.S. Patent No. 4,494,274; 584,370 which is a continuation of USSN No. 386,312, and 585,284, now U.S. Patent No. 5,034,413 which is a continuation of USSN 386,834, where the parent applications were filed on 27 July 1989. The disclosures of these patent applications are hereby expressly incorporated by reference.


Finally, certain EP₂-receptor agonists are disclosed in Nials et al, Cardiovascular Drug Reviews, Vol. 11, No. 2, pp. 165-179, Coleman et al, Comprehensive Medicinal Chemistry, Vol. 3, pp. 643-714, 1990 and Woodward et al, Prostaglandins, pp. 371-383, 1993.

Summary of the Invention

We have found that certain EP₂-receptor agonists are potent ocular hypotensive agents. We have further found that (+) trans-2-[-4(1-hydroxyhexyl) phenyl]-5-oxocyclopentaneheptanoic acid, and ester and unsaturated derivatives thereof, are especially useful in the treatment of glaucoma and surprisingly, cause no or significantly lower ocular surface hyperemia than the other compounds that are useful in lowering intraocular pressure, e.g. PGF₂ α and lower alkyl esters thereof.

The present invention relates to methods of treating ocular hypertension which comprises administering an effective amount of a compound represented by the formula I


5



wherein, the wavy bands indicate the α or β configuration, R is a saturated or unsaturated acyclic hydrocarbon group having from 1 to about 20 carbon atoms, or $\text{-(CH}_2\text{)}_m\text{R}_1$ wherein m is 0-10, and R_1 is an aliphatic ring having from about 3 to about 7 carbon atoms, or an aryl or heteroaryl ring having from about 4 to about 10 carbon atoms, e.g. R_1 may be cyclohexyl, phenyl, thienyl, pyridyl or furanyl, or a pharmaceutically acceptable salt thereof and the dashed bond represents either a single or double bond which may be in the cis or trans position. Preferably R_1 is lower alkyl.

More preferably the method of the present invention comprises administering a compound represented by the formula II

6



wherein R₂ is a lower alkyl radical and the other symbols are as defined above.

5

In a further aspect, the present invention relates to pharmaceutical compositions comprising a therapeutically effective amount of a compound of formulae (I) or (II) wherein the symbols have the above meanings, or a pharmaceutically acceptable salt thereof in admixture with a non-toxic, pharmaceutically acceptable liquid vehicle.


10

In a still further aspect, the present invention relates to certain (±) trans-2-[-4(1-hydroxyhexyl) phenyl]-5-oxocyclopentaneheptanoic acid, and ester and unsaturated derivatives thereof, of the above formulae, wherein the substituents and symbols are as defined hereinabove, or a pharmaceutically acceptable salt of such compounds.

15

Detailed Description of the Invention


5 The present invention relates to the use of (\pm) trans-2-[4(1-hydroxyhexyl) phenyl]-5-oxocyclopentaneheptanoic- acid, and ester and unsaturated derivatives thereof, as ocular hypotensives. These therapeutic agents are represented by compounds having the formula I



10


as defined above. The preferred compounds used in accordance with the present invention are encompassed by the following structural formula II

8



wherein R_2 is a lower alkyl radical.

5 In all of the above formulae, as well as in those provided hereinafter, the straight lines represent bonds. Where there is no symbol for the atoms between the bonds, the appropriate carbon-containing radical is to be inferred. For example in
10 formula I, the radical between the cyclopentyl ring and the



15 radical is a polymethylene (CH_2) radical, i.e. a hexylenyl radical. The dotted lines on the bond between carbons 5 and 6 (C-5), indicate a single or a double bond which can be in the cis or trans configuration. The radical adjacent the double
20 bond is a CH radical. If two solid lines are used that indicates a specific configuration for that double bond. Hatched lines at positions C-9 and C-11 indicate the α configuration. If one were to draw the β configuration, a solid triangular line would be used.

In the compounds used in accordance with the present invention, compounds having the C-9 or C-11 substituents in the α or β configuration are contemplated. As hereinabove mentioned, in all formulas provided herein broken line
5 attachments to the cyclopentane ring indicate substituents in the α configuration. Thickened solid line attachments to the cyclopentane ring indicate substituents in the β configuration.

For the purpose of this invention, unless further limited,
10 the term "alkyl" refers to alkyl groups having from one to ten carbon atoms and includes "lower alkyl" radicals having from one to five carbon atoms, the term "cycloalkyl" refers to cycloalkyl groups having from three to seven carbon atoms, the term "aryl" refers to aryl groups having from four to ten
15 carbon atoms. The term "saturated or unsaturated acyclic hydrocarbon group" is used to refer to straight or branched chain, saturated or unsaturated hydrocarbon groups having from one to about 6, preferably one to about 4 carbon atoms. Such groups include alkyl, alkenyl and alkynyl groups of
20 appropriate lengths, and preferably are alkyl, e.g. methyl, ethyl, propyl, butyl, pentyl, or hexyl, or an isomeric form thereof.

The definition of R may include a cyclic component,
25 $-(CH_2)_m R_1$, wherein m is 0-10, R_2 is an aliphatic ring from about 3 to about 7 carbon atoms, or an aromatic or heteroaromatic ring. The "aliphatic ring" may be saturated or unsaturated, and preferably is a saturated ring having 3-7 carbon atoms, inclusive. As an aromatic ring, R_1 preferably is
30 phenyl, and the heteroaromatic rings have oxygen, nitrogen or sulfur as a heteroatom, i.e., R_1 may be thienyl, furanyl, pyridyl, etc. Preferably m is 0-4.

Preferred representatives of the compounds within the
35 scope of the present invention are (\pm) trans-2-[-4(1-hydroxyhexyl)phenyl]-5-oxocyclopentaneheptanoic acid,

10

unsaturated derivatives thereof, and lower alkyl esters of these compounds.

5 The following compound may be used in the pharmaceutical compositions and the methods of treatment of the present invention are (\pm) trans-2-[-4(1-hydroxyhexyl) phenyl]-5-oxocyclopentaneheptanoic acid.

10 A pharmaceutically acceptable salt is any salt which retains the activity of the parent compound and does not impart any deleterious or undesirable effect on the subject to whom it is administered and in the context in which it is administered. Such salts are those formed with pharmaceutically acceptable cations, e.g., alkali metals, alkali
15 earth metals, etc.

Pharmaceutical compositions may be prepared by combining a therapeutically effective amount of at least one compound according to the present invention, or a
20 pharmaceutically acceptable salt thereof, as an active ingredient, with conventional ophthalmically acceptable pharmaceutical excipients, and by preparation of unit dosage forms suitable for topical ocular use. The therapeutically efficient amount typically is between about 0.0001 and about
25 5% (w/v), preferably about 0.001 to about 1.0% (w/v) in liquid formulations.

For ophthalmic application, preferably solutions are prepared using a physiological saline solution as a major
30 vehicle. The pH of such ophthalmic solutions should preferably be maintained between 4.5 and 8.0 with an appropriate buffer system, a neutral pH being preferred but not essential. The formulations may also contain conventional, pharmaceutically acceptable preservatives, stabilizers and
35 surfactants.

Preferred preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A preferred surfactant is, for example, Tween 80. Likewise, various preferred vehicles may be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose cyclodextrin and purified water.

Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

In a similar vein, an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated hydroxytoluene.

Other excipient components which may be included in the ophthalmic preparations are chelating agents. The preferred chelating agent is edentate disodium, although other chelating agents may also be used in place of or in conjunction with it.

12

The ingredients are usually used in the following amounts:

<u>Ingredient</u>	<u>Amount (% w/v)</u>
active ingredient	about 0.001-5
preservative	0-0.10
vehicle	0-40
tonicity adjustor	0-10
buffer	0.01-10
pH adjustor	q.s. pH 4.5-8.0
antioxidant	as needed
surfactant	as needed
purified water	as needed to make 100%

5 The actual dose of the active compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.

10 The ophthalmic formulations of the present invention are conveniently packaged in forms suitable for metered application, such as in containers equipped with a dropper, to facilitate application to the eye. Containers suitable for dropwise application are usually made of suitable inert, non-
15 toxic plastic material, and generally contain between about 0.5 and about 15 ml solution. One package may contain one or more unit doses.

20 Especially preservative-free solutions are often formulated in non-resealable containers containing up to about ten, preferably up to about five units doses, where a typical unit dose is from one to about 8 drops, preferably one to about 3 drops. The volume of one drop usually is about 20-35 μ l.

25 The invention is further illustrated by the following non-limiting Examples.

Example 1

(±) TRANS-2-[-4(1-HYDROXYHEXYL) PHENYL]-5-
OXOCYCLO-PENTANEHEPTANOICACID AND LOWER
ALKYL ESTERS THEREOF

The above acid compound is well known and may be
purchased or synthesized by methods known in the art. The
lower alkyl esters of this compound may be made by the
esterification procedures described in the various patent
applications described in the Background of the Invention.

Example 2**Intraocular Pressure**

Intraocular pressure was measured by
pneumatometry in normal monkeys. Studies were
performed in conscious animals trained to accept
pneumatometry. The compound of Example 1 was
administered topically to one eye in a 25 µl volume drop, the
contralateral eye received vehicle as a control. Statistical
analysis was by Student's paired t test.

The intraocular pressure results are summarized in
Table 1.

TABLE 1. EFFECT OF (\pm) TRANS-2-[-4(1-HYDROXYHEXYL) PHENYL]-5-OXOCYCLO PENTANE HEPTANOIC ACID, 0.1%, B.I.D. ON THE INTRAOCULAR PRESSURE OF NORMAL MONKEYS

	<u>TIME(HR) RELATIVE TO FIRST DOSE</u>	<u>NET CHANGE IN PRESSURE(MMHg)</u>
10	0	0
	2	-1.17
	4	-2.0
	6	-1.50
	24	-0.67
15	26	-1.0
	28	-1.83
	30	-1.50
	48	-1.50
	50	-2.33
20	52	-1.50
	54	-1.50
	72	-2.33
	74	-2.17
	76	-2.33
25	78	-2.0
	96	-1.83
	98	-2.0
	100	-2.0
30	102	-2.0

(all changes in intraocular pressure are significant $p < 0.01$ according to Students t test)

Example 3

35 Intraocular Pressure Reduction After Laser Treatment

Intraocular pressure reduction was also achieved in laser-induced ocular hypertensive monkeys. Ocular hypertension was induced by photocoagulating the trabecular meshwork by circumferential argon laser treatment.

40

45

TABLE 2. EFFECT OF A SINGLE DOSE (0.1%) OF TRANS-2-[-4(1-HYDROXYHEXYL)PHENYL]-5-OXO-CYCLOPENTANEHEPTANOIC ACID ON OCULAR PRESSURE ON LASER-INDUCED OCULAR HYPERTENSIVE MONKEYS

	<u>TIME (HR) RELATIVE TO FIRST DOSE</u>	<u>CHANGE IN INTRA- OCULAR PRESSURE (MMHg)</u>
10	0	0
	1	-4.5**
	2	-6.0**
	4	-4.7**
	6	-4.5*

* $p < 0.05$

** $p < 0.01$ (Students t test)

Example 4


Determination of EP₂ Receptor Activity

EP₂ receptor activity may be measured in accordance with the procedure disclosed in Woodward et al, Prostaglandins, pp. 371-383, 1993, which is hereby incorporated by reference in its entirety.

The foregoing description details specific methods and compositions that can be employed to practice the present invention, and represents the best mode contemplated. However, it is apparent from one of ordinary skill in the art that different pharmaceutical compositions may be prepared and used with substantially the same results. That is other EP₂-receptor agonists, such as 19R(OH)PGE₂, butaprost, etc. will effectively lower intraocular pressure in animals and are within the broad scope of the present invention.

Claims:


1. A method of treating ocular hypertension which comprises applying to the eye an amount sufficient to treat ocular hypertension of a compound of formula I



- wherein the wavy bonds indicate that α or β configuration, the dashed bond represents either a single or double bond, which double bond may be a cis or trans double bond and R is a saturated or unsaturated acyclic hydrocarbon group having from 1 to about 20 carbon atoms, or $-(CH_2)_mR_1$ wherein m is 0-10, and R_1 is an aliphatic ring having from about 3 to about 7 carbon atoms, or an aryl or heteroaryl ring having from about 4 to about 10 carbon atoms and wherein the heteroatom is selected from the group consisting of N, O and S.

2. The method of Claim 1 wherein said compound is a prostaglandin derivative of the formula I

17




wherein R₂ is a lower alkyl radical, the hatched line indicates the α configuration and the solid triangular line indicates the β configuration.

3. The method of Claim 2 wherein said compound is (±) trans-2-[-4(1-hydroxyhexyl) phenyl]-5-oxocyclopentane-heptanoicacid.

4. A pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I


18



5 wherein the wavy bonds indicate that α or β configuration, the dashed bond represents either a single or double bond, which double bond may be a cis or trans double bond and R is a saturated or unsaturated acyclic hydrocarbon group having from 1 to about 20 carbon atoms, or $-(CH_2)_mR_1$ wherein m is 0-10, and R_1 is an aliphatic ring having from about 3 to about 7 carbon atoms, or an aryl or heteroaryl ring having from about 4 to about 10 carbon atoms and wherein the heteroatom is selected from the group consisting of N, O and S.

15 5. An ophthalmic solution comprising a therapeutically effective amount of a compound of formula I

19




5 wherein the wavy bonds indicate that α or β configuration, the
dashed bond represents either a single or double bond, which
double bond may be a cis or trans double bond and R is a
saturated or unsaturated acyclic hydrocarbon group having
from 1 to about 20 carbon atoms, or $-(CH_2)_mR_1$ wherein m is
0-10, and R_1 is an aliphatic ring having from about 3 to about
7 carbon atoms, or an aryl or heteroaryl ring having from
10 about 4 to about 10 carbon atoms and wherein the heteroatom
is selected from the group consisting of N, O and S.

15 6 The ophthalmic solution of Claim 5 comprising at
least one ingredient selected from the group of an
ophthalmically acceptable preservative, buffer system,
antioxidant and chelating agent.

20 7. The ophthalmic solution of Claim 6 wherein said
compound is a prostaglandin derivative of the formula I

20



wherein R_2 is a lower alkyl radical, the hatched line indicates the α configuration and the solid triangular line indicates the β configuration.


8. The ophthalmic solution of Claim 7 wherein said compound is (\pm) trans-2-[-4(1-hydroxyhexyl) phenyl]-5-oxocyclopentaneheptanoicacid, and ester and unsaturated derivatives thereof.

9. A pharmaceutical product, comprising
a container adapted to dispense its contents in metered form; and
an ophthalmic solution therein, as defined in Claim 5.

20

10. A compound of the formula I

21



5 wherein the wavy bonds indicate that α or β configuration, the dashed bond represents either a single or double bond, which double bond may be a cis or trans double bond and R is a saturated or unsaturated acyclic hydrocarbon group having from 1 to about 20 carbon atoms, or $-(\text{CH}_2)_m\text{R}_1$ wherein m is 0-10, and R_1 is an aliphatic ring having from about 3 to about 7 carbon atoms, or an aryl or heteroaryl ring having from about 4 to about 10 carbon atoms and wherein the heteroatom is selected from the group consisting of N, O and S.

11. A method for treating intraocular pressure which comprises applying to the eye an amount sufficient to treat ocular hypertension of a compound having EP₂ receptor agonist activity.

INTERNATIONAL SEARCH REPORT

In International Application No

PCT/US 95/00288

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07C405/00 A61K31/557

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF OCULAR PHARMACOLOGY, vol. 10, no. 1, 1994 pages 177-193, D.F. WOODWARD ET AL. 'Studies on the ocular hypotensive effects of prostaglandin F ₂ alpha ester prodrugs and receptor selective prostaglandin analogs' see page 177 see page 180, paragraph 2 -----	1-11

☐ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

17 May 1995

Date of mailing of the international search report

06.06.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Tzschoppe, D